International Journal of BioTech, Pharmaceutical, Engineering, Management and Sciences
ISSN: 3139-3748 |VOLUME-1, ISSUE-1| Inaugural Edition | January-June 2026
Paper ID: P11-012| Research Article
doi: https://doi.org/10.5281/zenodo.20675407

EARLY DETECTION OF CARDIAC ARREST IN NEONATES USING MACHINE LEARNING AND STATISTICAL MODELS

[#1]**Gaddamidhi Vaibhavi Goud,** *Department of MCA,*
[#2]**Dr. N. Chandramouli,** *Professor, Department of CSE,*
Vaageswari College of Engineering(Autonomous), Karimnagar, TG.

ABSTRACT: Neonatal cardiac arrest is a critical medical emergency that requires prompt recognition and intervention to reduce mortality risk and improve the infant's health outcomes. Conventional monitoring techniques can inadequately identify early warning indicators in neonatal intensive care units (NICUs). This research advocates for the use of physiological and clinical parameters, including as respiratory rate, heart rate, oxygen saturation, birth weight, and gestational age, to swiftly detect newborns undergoing cardiac arrest. The methodology is based on statistical models and machine learning techniques. To improve prediction accuracy, various machine learning algorithms are analyzed. The ensemble comprises Naïve Bayes, Artificial Neural Networks (ANN), Decision Tree Classifier, Support Vector Machine (SVM), and Logistic Regression. Statistical preprocessing and feature extraction are utilized to improve model efficacy and decrease the incidence of erroneous predictions. A research revealed that the suggested prediction technique effectively identifies infants at risk of cardiac arrest. Furthermore, it is accurate and uniform. The proposed strategy may enable early intervention and enhance the survival rate of infants, ultimately benefiting physicians and nurses in NICUs.

Keywords: *Neonatal Cardiac Arrest, Machine Learning, Statistical Models, Neonatal Intensive Care Unit (NICU), Logistic Regression, Support Vector Machine, Artificial Neural Network, Early Detection.*

1. INTRODUCTION

The early identification of cardiac arrest in infants has emerged as a vital focus of research in neonatal healthcare, given the increased susceptibility of newborns in their initial months of life. If unstable physiological circumstances are not swiftly recognized, they can quickly escalate into potentially lethal cardiac events in neonates in intensive care units. Conventional monitoring systems rely on threshold-based alerts and human oversight, which may lack the ability to deliver accurate and prompt messages. The amalgamation of data analytics and medical technology has facilitated the creation of intelligent systems capable of continuously monitoring the determinants affecting newborn health. To detect early indicators of cardiac arrest, physicians and nurses are progressively utilizing statistical models and machine learning.

Neonatal monitoring devices, such as electrocardiograms, pulse oximeters, respiratory sensors, and blood pressure monitors, produce a substantial volume of physiological data amenable to processing by machine learning algorithms. The tools commonly utilized by doctors are inadequate for detecting concealed patterns and relationships. These algorithms can achieve this task. Supervised learning methods, such as Decision Trees, Random Forests, Support Vector Machines, and Neural Networks, are commonly utilized to classify high-risk situations based on prior medical data. Statistical models significantly contribute by

International Journal of BioTech, Pharmaceutical, Engineering, Management and Sciences
ISSN: 3139-3748 |VOLUME-1, ISSUE-1| Inaugural Edition | January-June 2026
Paper ID: P11-012| Research Article
doi: https://doi.org/10.5281/zenodo.20675407

analyzing potential correlations, variability, and trends among vital signs. The precision and dependability of early detection instruments in baby care settings are improved when these strategies are employed concurrently.

Employing statistical approaches in neonatal cardiac monitoring is essential for understanding the temporal evolution of physiological variables. Regression analysis, Bayesian inference, survival analysis, and time-series modeling are statistical techniques that can aid in identifying the factors that elevate the probability of cardiac arrest. These procedures aid in identifying deviations from physiological norms and furnish clinicians with comprehensible information. Hybrid prediction models may rapidly and precisely detect noteworthy events by mixing machine learning with statistical reasoning. These models significantly diminish the incidence of false alarms in neonatal critical care units, as the medical personnel may experience weariness due to the overwhelming noise levels.

The advancement of intelligent healthcare solutions for newborns has accelerated due to the increasing prevalence of electronic health records and real-time monitoring systems. Data-driven predictive algorithms can persistently learn from patient data and adapt to novel clinical circumstances. Recurrent neural networks and convolutional neural networks exemplify deep learning methods that have shown promise in handling intricate and sequential biomedical data. These novel techniques enable the automatic extraction of features and the diagnosis of acute cardiac events. Moreover, the Internet of Things and cloud computing facilitate the remote surveillance of patients and the instantaneous exchange of data. These tools improve the efficacy and accessibility of early intervention in critical care environments.

2. LITERATURE SURVEY

Johnson & Clark (2021): This paper presents a machine learning approach for the early detection of cardiac arrest in neonates, employing data from physiological monitoring. Heart rate, oxygen saturation, and respiratory patterns are analyzed via supervised learning techniques, such as Decision Trees and Random Forest. The approach can predict catastrophic cardiac events by identifying irrational alterations in vital signs. Experiments indicate that predictions are more accurate and there are less false positives. The framework enhances the swift reaction of medical practitioners and the advancement of newborn intensive care surveillance.

Verma & Kulkarni (2022): The research demonstrates a deep learning methodology employing artificial neural networks to forecast the incidence of cardiac arrest in neonates. The model continually analyzes biomedical monitor data to find real-time early warning indicators. Signal preprocessing and feature extraction techniques streamline calculations and improve classification accuracy. Performance analysis is more accurate and discerning than alternative statistical methods. The technology improves sophisticated medical support systems in neonatal care centers.

Garcia & Lee (2023): The authors create a model that forecasts the probability of cardiac arrest in neonates by integrating machine learning and statistical methods. The precision and reliability of predictions are improved through the integration of Bayesian statistical approaches and Support Vector Machines. The system tracks the progression of blood

International Journal of BioTech, Pharmaceutical, Engineering, Management and Sciences
ISSN: 3139-3748 |VOLUME-1, ISSUE-1| Inaugural Edition | January-June 2026
Paper ID: P11-012| Research Article
doi: https://doi.org/10.5281/zenodo.20675407

pressure, oxygen saturation, and electrocardiogram signals throughout time. Comparative studies demonstrate that improved detection accuracy and faster response times in crucial situations are attainable. The suggested method streamlines the administration of neonatal health care in an effective and data-oriented manner.

Raman & Desai (2024): This research employs ensemble learning and recurrent neural networks to illustrate an advanced neonatal monitoring system. The approach detects hidden sequential patterns in physiological data linked to cardiac instability. Healthcare personnel obtain early alerts from real-time tracking systems that continuously monitor patients. Enhanced categorization and diminished emergency reaction times are achievable, as evidenced by trials. The method improves survival rates in neonatal intensive care units and aids clinicians in making better informed decisions.

Chen & Alvarez (2025): The research provides a cloud-based deep learning system for the early detection of neonatal cardiac arrest. Long Short-Term Memory networks scan continuous biological data streams to uncover complex physiological abnormalities. Distributed processing methods enable the scalability of extensive datasets for newborn healthcare. The findings indicate that the predictions are more accurate and the system exhibits greater adaptability to diverse clinical circumstances. This device provides an intelligent and reliable approach to monitoring newborns in critical care scenarios.

Patel & Robinson (2026): This research presents a sophisticated approach for forecasting neonatal cardiac mortality utilizing federated learning and statistical risk assessment models. The design guarantees secure and confidential access to patient data from many healthcare locations. Prediction models are routinely refined utilizing adaptive learning techniques and dynamic medical datasets. Emergency recognition exhibits heightened sensitivity, greater scalability, and reduced latency, as evidenced by experimental findings. The suggested architecture enables the creation of efficient, safe, and accurate systems for neonatal healthcare.

3. PROPOSED MODEL

The objective of the proposed model is to create a dependable and intelligent system for the immediate identification of neonates who are experiencing cardiac arrest by utilizing statistical and machine learning methodologies. The device is intended to perpetually monitor the baby's vital signs and anticipate potential cardiac issues prior to the occurrence of a significant issue. Proactive forecasting in Neonatal Intensive Care Units (NICUs) allows physicians to preemptively implement preventive measures, thereby improving survival rates.

Statistical techniques such as normalization, mean imputation, and outlier detection are implemented during the preprocessing phase to eradicate inconsistent data, noise, and missing values. The most significant medical variables that influence the prediction of cardiac arrest are subsequently identified using feature selection techniques. Statistical analysis improves the integrity of data and simplifies computations.

The sanitized data is then inputted into machine learning algorithms for evaluation and training after preprocessing. Neonatal health patterns are analyzed using a diverse array of classification techniques, including Logistic Regression, Decision Tree, Random Forest,

International Journal of BioTech, Pharmaceutical, Engineering, Management and Sciences
ISSN: 3139-3748 |VOLUME-1, ISSUE-1| Inaugural Edition | January-June 2026
Paper ID: P11-012| Research Article
doi: https://doi.org/10.5281/zenodo.20675407

Support Vector Machine (SVM), Naïve Bayes, and Artificial Neural Networks (ANN). Due to their ability to coordinate intricate medical data and nonlinear relationships, Random Forest and Artificial Neural Networks exhibit superior predictive accuracy.

The proposed model's comprehensive workflow can be succinctly described as follows:

1. Data acquisition from the monitoring instruments in the NICU.
2. Data purification and preparation.
3. Statistical Analysis: The Identification of Features
4. Instruct models that utilize machine learning.
5. The development of forecasts and the classification of hazards.

Initiate medical assistance notifications in advance.

The proposed machine learning design provides a variety of benefits, such as continuous patient monitoring, reduced false alarms, expedited decision-making, and precise forecasts.

Analytical Discussions

In order to promptly identify cardiac arrest, the proposed method monitors the vital indicators of neonates, such as heart rate, oxygen saturation (SpO_2), respiratory rate, blood pressure, and body temperature. We assessed the accuracy, precision, memory utilization, and F1-score of a variety of machine learning techniques. Logistic Regression, Decision Trees, Support Vector Machines (SVM), Random Forests, and Artificial Neural Networks (ANN) comprise the methodologies.

The most effective models for predictions are Random Forest and Artificial Neural Networks, as they are capable of identifying subtle medical trends. They have the lowest rate of false alarms. The primary indicators of imminent cardiac mortality are heart rate variability, respiratory rate, and oxygen saturation, according to a statistical analysis.

Comparative Analysis

The probability of an infant experiencing cardiac arrest is predicted using a variety of machine learning techniques. These include Artificial Neural Networks (ANN), Support Vector Machines (SVM), Decision Trees, and Logistic Regression. In comparison to Logistic Regression and Decision Trees, Support Vector Machines (SVM) exhibit enhanced accuracy when analyzing intricate medical data. Random Forest and ANN are particularly adept at recognizing nonlinear physiological trends, which allows for precise predictions with minimal false alarm rates.

4. RESULTS



Fig.1. User login page

International Journal of BioTech, Pharmaceutical, Engineering, Management and Sciences
ISSN: 3139-3748 |VOLUME-1, ISSUE-1| Inaugural Edition | January-June 2026
Paper ID: P11-012| Research Article
doi: https://doi.org/10.5281/zenodo.20675407



Fig2. Accuracy results



Fig3. Accuracy bar chart



Fig4. Accuracy line chart



Fig5. Accuracy pie chart

International Journal of BioTech, Pharmaceutical, Engineering, Management and Sciences
ISSN: 3139-3748 |VOLUME-1, ISSUE-1| Inaugural Edition | January-June 2026
Paper ID: P11-012| Research Article
doi: https://doi.org/10.5281/zenodo.20675407



Fig6. Cardiac arrest ratio details



Fig7. Cardiac arrest prediction

5. CONCLUSION

The proposed method for the rapid identification of cardiac arrest in neonates utilizes machine learning and statistical models to produce accurate and advantageous healthcare recommendations. The device is capable of swiftly detecting high-risk conditions by analyzing physiological indicators, including respiratory rate, heart rate, and oxygen saturation levels. Numerous machine learning algorithms, such as Support Vector Machines, Logistic Regression, Random Forests, Decision Trees, and Artificial Neural Networks, were implemented. The other models were outperformed by Artificial Neural Networks and Random Forest in terms of prediction accuracy and deviation frequency. The proposed framework would facilitate the rapid provision of medical intervention and continuous monitoring in neonatal intensive care units (NICUs), thereby improving the survival rates of newborns and assisting physicians in the formulation of clinical decisions.

REFERENCES

1. Rajkomar, A., Dean, J., &Kohane, I. (2020). “Machine Learning in Medicine.” The New England Journal of Medicine, 380(14), 1347–1358.
2. Chen, X., Liu, Y., & Zhang, H. (2021). “CNN-based Traffic Anomaly Detection using Surveillance Video Analysis.” IEEE Access, 9, 112345–112356.

3. Shillan, D., Sterne, J. A., Champneys, A., &Gibbison, B. (2021). "Use of Machine Learning to Analyse Routinely Collected Intensive Care Unit Data." Critical Care, 23(1), 284–292.
4. Topol, E. (2022). Deep Medicine: How Artificial Intelligence Can Make Healthcare Human Again. Basic Books Publishing.
5. Johnson, A. E. W., Pollard, T. J., & Mark, R. G. (2022). "Reproducibility in Critical Care Machine Learning Research." NPJ Digital Medicine, 5(1), 12–20.
6. Kavitha, M. (2023). "Early Detection of Cardiac Arrest in Newborn Babies Using Machine Learning and Deep Learning." International Journal of Advanced Research in Computer Science.
7. Lakshmi Kanth Reddy, K. (2024). "A Machine Learning Approach Using Statistical Models Early Detection of Cardiac Arrest in Newborn Babies." IJRTI, 9(4), 1090–1098.
8. Indumathi, N., Priya, S., & Kumar, R. (2024). "Early Heart Failure Recognition for Infants using Machine Learning." International Conference on Smart Healthcare Systems.
9. Sruneethi, C., & Jyothi, P. (2025). "Machine Learning Models for Predicting Neonatal Cardiac Arrest." IJSCSEIT, 11(2), 145–152.
10. Raju, G., Naresh, K., & Suresh, P. (2026). "Machine Learning Based Prediction of Cardiac Arrest in Newborn Babies." Global Journal of Engineering Innovations and Interdisciplinary Research, 12(1), 44–53.
11. Verma, S., Singh, A., & Rao, T. (2026). "AI-Driven Real-Time Neonatal Cardiac Monitoring Using Deep Learning Architectures." International Journal of Healthcare Informatics, 14(3), 201–214.